

Fokker-Planck equation of distributions of financial returns and power laws

Didier Sornette^{1,2}

¹ *Laboratoire de Physique de la Matière Condensée, CNRS UMR6632*

Université des Sciences, B.P. 70, Parc Valrose, 06108 Nice Cedex 2, France

² *Department of Earth and Space Science*

and Institute of Geophysics and Planetary Physics

University of California, Los Angeles, California 90095, USA

Abstract: Our purpose is to relate the Fokker-Planck formalism proposed by [Friedrich et al., Phys. Rev. Lett. 84, 5224 (2000)] for the distribution of stock market returns to the empirically well-established power law distribution with an exponent in the range 3 – 5. We show how to use Friedrich et al.’s formalism to predict that the distribution of returns is indeed asymptotically a power law with an exponent μ that can be determined from the Kramers-Moyal coefficients determined by Friedrich et al. However, with their values determined for the U.S. dollar-German mark exchange rates, the exponent μ predicted from their theory is found around 12, in disagreement with the often-quoted value between 3 and 5. This could be explained by the fact that the large asymptotic value of 12 does not apply to real data that lie still far from the stationary state of the Fokker-Planck description. Another possibility is that power laws are inadequate. The mechanism for the power law is based on the presence of multiplicative noise across time-scales, which is different from the multiplicative noise at fixed time-scales implicit in the ARCH models developed in the Finance literature.

I. SUMMARY OF COMPETING MODELS FOR RETURN DISTRIBUTIONS

The distribution of asset returns provides the zeroth-order description of the statistical properties of asset returns. Over the years, several competing models have been proposed to describe the non-Gaussian heavy-tail structure of asset returns. Without being exhaustive, we can cite:

- auto-regressive conditional heteroskedastic (ARCH) [1] models which assume that the variance is a function of past price variations and possibly of past variances (GARCH) [2,3]; these models lead in general to power tails;
- truncated Lévy laws [4,5];
- multiplicative noise models with reinjections [6–8] which provide a mechanism for power law distributions [9–13]; we note that such models are equivalent to ARCH models via a nonlinear change of variable [14,15,8];
- stretched exponential models [16,17] which can be derived as tails of multiplicative processes [18];
- multifractal cascade models from large to small time-scales [19–21].

In a recent letter [22], Friedrich et al. have proposed an alternative approach in terms of a Fokker-Planck equation for the distribution of asset returns at different time-scales, relying on their previous similar Fokker-Planck approach for the description of the distribution of velocity increments in turbulent cascades [23] and their proposed analogy between turbulent cascades in hydrodynamics and information cascades in stock markets.

Defining the price increment

$$\Delta x(t, \Delta t) = x(t + \Delta t) - x(t) \tag{1}$$

over the time interval Δt (which also defines the time-scale), the purpose of Friedrich et al. [22] is to write down an equation for the statistical process underlying the price changes

Δx over a series of nested time delays Δt_i (or time-scales) of decreasing durations. The controlling parameter is thus Δt while t is dummy variable which, when varied, gives different realizations of the same process at the given time-scale Δt . In other words, the problem is to write down an equation of $\Delta x(\Delta t)$ as a function of time-scale Δt . In this goal, using the insight provided by cascade models, it is natural to define the logarithmic time-scale

$$\tau = \ln(t_0/\Delta t) , \quad (2)$$

where $t_0 = 40960s$ is the large time-scale from which the cascade proceeds. Friedrich et al. [22] then arrive at the following Langevin equation for the price increment $\Delta x(\tau)$ as a function of logarithmic time-scale τ

$$d\Delta x(\tau) = D^{(1)}(\Delta x) d\tau + \sqrt{D^{(2)}(\Delta x, \tau)} dW , \quad (3)$$

where the Kramers-Moyal coefficients are given by

$$D^{(1)}(\Delta x) = -0.44 \Delta x , \quad (4)$$

$$D^{(2)}(\Delta x, \tau) = 0.003e^{-\tau/2} + 0.019(\Delta x + 0.04)^2 , \quad (5)$$

for the U.S. dollar-German mark exchange rates. dW is the increment of the random walk $W(\tau)$, with white noise spectrum and variance equal to 2. This Langevin equation (3) is equivalent to the Fokker-Planck equation describing the distribution $P(\Delta x, \tau)$ of price variations Δx at a given logarithmic time-scale τ [22]:

$$\frac{\partial P(\Delta x, \tau)}{\partial \tau} = -\frac{\partial D^{(1)}(\Delta x)P(\Delta x, \tau)}{\partial \Delta x} + \frac{\partial^2 D^{(2)}(\Delta x)P(\Delta x, \tau)}{\partial \Delta x^2} . \quad (6)$$

For a fixed τ , i.e., time-scale Δt , $P(\Delta x, \tau)$ is nothing but the usual distribution of price increments studied in many empirical works. The innovation brought by the formulation (6) or its equivalent stochastic version (3) is to relate the distribution at one time-scale to that at other time-scales. This formulation expresses in a familiar framework (the Fokker-Planck or Langevin equations) the cascade models already discussed in Refs. [19–21]. Let us stress again that the “time” τ entering in (6) and (3) is a time-scale, not a time along which one

follows the trajectory of the price or of the price increments. Thus, equations (6) and (3) are not equations of the dynamical evolution of prices. They are mathematical representations of the cascade model [19–21]. In particular, equation (3) is a stochastic equation describing how price increments change with time-scale. For a single realization of the noise trajectory $W(t)$, it describes one particular cascade path of the price increments as a function of time-scale. Constructing an ensemble of noise trajectories flowing from the large time-scale t_0 down to some fixed time-scale Δt will then provide an ensemble of price variations at this time-scale Δt from which the distribution at this time-scale can be determined. We now turn to examine how (3) can be used precisely in this goal of calculating $p(\Delta x, \tau)$.

II. MAPPING OF THE LANGEVIN FORMALISM OF FRIEDRICH ET AL. ONTO INTERMITTENTLY AMPLIFYING MULTIPLICATIVE NOISE

A. Multiplicative noise with reinjection

In all the derivation below, we thus consider a logarithmic time-scale τ flowing from 0 (large time-scale $\Delta t = t_0$) to a fixed value τ corresponding to a fixed time-scale Δt . Our goal is to consider many trajectories of equal logarithmic time-scale duration τ and derive the corresponding distribution of price increments Δx at this fixed time scale Δ .

To perform this mapping, the key remark is that, for large Δx , $\sqrt{D^{(2)}(\Delta x, \tau)}$ reduces to approximately $\sqrt{0.019} \Delta x$, which corresponds to a multiplicative noise. In contrast, when Δx becomes small, $\sqrt{D^{(2)}(\Delta x, \tau)}$ reduces to a function independent of Δx and the multiplicative noise is transformed into an additive noise.

These two limiting behaviors, when taken together, lead exactly to the mechanism of multiplicative noise with reinjection, shown to generate power law distribution, discussed in Ref. [6]. Specifically, discretizing (3) by considering a small time-scale increment $d\tau$, we get

$$\Delta x(\tau + d\tau) = a(\tau)\Delta x(\tau) , \quad \text{for large } \Delta x , \quad (7)$$

where

$$a(\tau) = 1 - 0.44d\tau + \sqrt{0.019} dW . \quad (8)$$

Note that $a(\tau)$ becomes larger than 1 when $-0.44d\tau + \sqrt{0.019} dW$ is positive, corresponding to an amplification or growth of $\Delta x(\tau)$. In constrast, for small Δx , the expression (7) is changed into

$$\Delta x(\tau + d\tau) = a_0(\tau)\Delta x(\tau) + \sigma dW , \quad \text{for small } \Delta x , \quad (9)$$

where $a_0 = 1 - 0.44d\tau$ and

$$\sigma = 0.003e^{-\tau/2} + 0.019(0.04)^2 . \quad (10)$$

Without the time-scale dependence of σ , expression (9) would be nothing but the standard mean-reversal equation (albeit in time-scale rather than time) or Orstein-Uhlenbeck process.

B. Properties of the fluctuations of $\Delta x(\tau)$

The stochastic dynamics described by (7) and (9) is the following. For most of the time-scales, the realizations of the random noise dW are small and the multiplicative coefficient $a(\tau)$ is smaller than 1. As a consequence, Δx shrinks and eventually the additive noise equation (9) takes over and ensures that Δx does not go to zero. However, due to the intrinsic stochastic nature of the noise dW , there will be random occurrences of dW that make the multiplicative factor $a(\tau)$ larger than 1. When this occurs, Δx is amplified. If this occurs over several successive time-scale steps, Δx is exponentially amplified. This is the mechanism that leads to heavy tailed distribution, by intermittent multiplicative amplification. Such amplification is bound to have a finite lifetime since the expectation of the multiplicative growth rate $\langle \ln a(\tau) \rangle = a_0 = 1 - 0.44d\tau$ is negative.

We can make this argument more precise by quantifying the power law distribution $P(\Delta x)$ generated by the mechanism involving the “fight between exponentials” [26]. As we have just said, large Δx are generated by intermittent amplifications resulting from the

multiplication by several successive values of a larger than one. We now give a mean-field argument to derive the result that this process produces distributions with a power law tail.

We present the argument using a discretization of the time-scale in “unit” steps $d\tau$. Let us call $p_>$ the probability that the multiplicative factor a is found larger than 1. The probability to observe n successive multiplicative factors a larger than 1 over n successive time-scale steps $d\tau$ is thus $p_>^n$. Let us call $a_>$ the average of a conditioned on being larger than 1: $a_>$ is thus the typical value of the amplification factor. When n successive multiplicative factors occur with values larger than 1, they typically lead to an amplification of the amplitude of Δx by $a_>^n$. Using the fact that the equation (9) ensures that the amplitude of Δx remains of the order of σ when the multiplicative factors a are less than 1, this shows that a value of Δx of the order of $\Delta x \approx \sigma a_>^n$ occurs with probability

$$p_>^n = \exp(n \ln p_>) \approx \exp\left(\ln p_> \frac{\ln \frac{\Delta x}{\sigma}}{\ln a_>}\right) = \frac{1}{(\Delta x/\sigma)^\mu} \quad (11)$$

with $\mu = -\ln p_>/\ln a_>$, i.e.

$$p_> a_>^\mu = 1 . \quad (12)$$

This last expression represents a kind of mean field version of the exact solvability condition [24,14,25,6,8]

$$\langle a^\mu \rangle = 1 , \quad (13)$$

determining the exponent μ of the power law tail of the distribution of Δx :

$$P(\Delta x) \sim \Delta x^{-(1+\mu)} . \quad (14)$$

The power law distribution is thus the result of an exponentially small probability of creating an exponentially large value. Expression (11) does not provide a precise determination of the exponent μ , only an approximate one since we have used a kind of mean-field argument in the definition of $a_>$. It however illuminates the physical mechanism for the power law, as resulting from a fight between exponentials [26].

The exact value of the exponent is instead determined by the non-mean field condition (13) which is derived as follows. In the presence of multiplicative noise, the distribution $P(\Delta x)$ is solution of the integral equation which, for large Δx , reads

$$P(\Delta x) = \int \frac{da}{a} P_a(a) \int d\Delta x P(\Delta x/a) , \quad (15)$$

where $P_a(a)$ is the distribution of the multiplicative factors defined in (7). Looking for a power law solution (14) put in (15) gives directly the equation (13) on the exponent μ . Our somewhat heuristic summary presented here is backed up by exact rigorous analysis [24].

C. Numerical evaluation

From expression (8), we see that $a(\tau)$ is Gaussian. For small time-scale increments $d\tau$, $\ln a(\tau) = -0.44d\tau + \sqrt{0.019} dW$ and we do not need to distinguish between normality and lognormality, and we have

$$\langle \ln a \rangle = -0.44d\tau \quad (16)$$

$$\langle (\ln a)^2 \rangle - \langle \ln a \rangle^2 = 2 \times 0.019 d\tau, \quad (17)$$

where we have used the property of the Wiener process $\langle (dW)^2 \rangle = 2d\tau$ (we keep the same convention as in Friedrich et al. [22] of a variance equal to 2). This allows us to use the exact solution of (13) for the exponent μ [6]:

$$\mu = - \frac{\langle \ln a \rangle}{\langle (\ln a)^2 \rangle - \langle \ln a \rangle^2} . \quad (18)$$

The $d\tau$ dependence cancels out between the numerator and denominator, and with the value obtained for the UD dollar-German Mark exchange rates, we get the estimation $\mu \approx 11.6$. This value is significantly larger than the value in the range 3 – 5 often reported in the literature [9–13] for assets others than exchange rates. One possible explanation is that exchange rates are known to exhibit thinner tails than stocks. Another explanation is that the parameters of the distributions within the Fokker-Planck formalism are far from their

stationary values and the value of the exponent in the range $3-5$ could be a non-asymptotic value. Another explanation is that the description by power laws, even if it has a rather long history, is not adequate [16].

III. CONCLUSION

In summary, we have used the stochastic Langevin formalism associated with the Fokker-Planck description offered by Friedrich et al. [22] to show that, in their model, the asymptotic tail of the distribution of price variations at a given time-scale is a power law with a large exponent $\mu \approx 11.6$. We have stressed that the stochastic Langevin equation or equivalently the Fokker-Planck formalism are expressed in terms of an evolution along the time-scale axis rather than along the time arrow. The power law distribution obtained in this model thus relies intrinsically in the assumed cascade from large time-scales to small time-scales and is thus different from the multiplicative models that are equivalent to usual ARCH formulations [14,15,8], since the later are defined for a fixed time scale.

However, since the multiplicative cascade across time-scales of Friedrich et al. as well as multiplicative noises at a fixed time scale can both produce fat tail power law distributions, the empirical observation of such fat tails and their adequate fits by this formalism [22] is insufficient to prove the reality of the cascade. In order to demonstrate the existence of a genuine cascade across scales, it is necessary to calculate in addition the correlation functions between different time-scales and at varying time lags: the asymmetry of the correlation functions across time-scales reported in [20,21] is, in our opinion, a convincing way of proving the existence of the cascade. This was also noted independently by the Olsen group in Zurich for the foreign exchange market, who coined it the HARCH effect [27,3]: the coarse-grained volatility predicts the fine-grained volatility better than the other way around. They also found this effect for the implied forward rates derived from Eurofutures contracts [28].

REFERENCES

- [1] R.F. Engle, *Econometrica* 50, 987 (1982).
- [2] T. Bollerslev, R.Y. Chous and K.F. Kroner, ARCH modeling in Finance - A review of the theory and empirical evidence, *J. Econometrics* 52, 5-59 (1992).
- [3] M.M. Dacorogna, U.A. Mller, R.B. Olsen, O.V. Pictet, Modelling Short-Term Volatility with GARCH and HARCH Models, in “Nonlinear Modelling of High Frequency Financial Time Series,” by C. Dunis, B. Zhou (John Wiley & Sons, 1998).
- [4] R.N. Mantegna and H.E. Stanley, *Nature* 376, 46 (1995); I. Koponen, *Phys. Rev.* 52, 1197 (1995).
- [5] A. Arnéodo, J.-P. Bouchaud, R. Cont, J.-F. Muzy, M. Potters and D. Sornette, Comment on ”Turbulent cascades in foreign exchange markets”, preprint at cond-mat/9607120.
- [6] D. Sornette and R. Cont, *J. Phys. I France* **7**, 431 (1997);
- [7] H. Takayasu, A.-H. Sato and M. Takayasu, *Phys. Rev. Lett.* **79**, 966 (1997); D. Sornette, *Phys. Rev. E* 57, 4811 (1998)
- [8] D. Sornette, *Physica A* 250, 295 (1998).
- [9] D.G. de Vries, Stylized Facts of Nominal Exchange Rate Returns, S. 348 - 89 in: van der Ploeg, F., ed., *The Handbook of International Macroeconomics*. Blackwell: Oxford (1994).
- [10] A. Pagan, *Journal of Empirical Finance* 3, 15 - 102 (1996).
- [11] D.M. Guillaume, Dacorogna, M.M., Davé, R.R, Mller, J.A., Olsen, R.B. & Pictet, O.V., *Finance & Stochastics* 1, 95 - 129 (1997).
- [12] P. Gopikrishnan, Meyer, M., Amaral, L.A.N. & Stanley, H.E., *European Physical Journal B* 3, 139 - 140 (1998); P. Gopikrishnan, V. Plerou, X. Gabaix and H.E. Stan-

- ley, Statistical Properties of Share Volume Traded in Financial Markets (preprint at <http://arXiv.org/abs/cond-mat/0008113>)
- [13] E. Jondeau and M. Rockinger, The Tail Behavior of Stock Returns: Emerging versus Mature Markets. mimeo: HEC - School of Management (1999).
 - [14] de Haan, L., Resnick, S.I., Rootzén, H. & de Vries, C.G., Stochastic Processes and Applics. **32**, 213 (1989).
 - [15] Embrechtz, P., Kluppelberg C. and Mikosch T., *Modelling extremal events*, Springer, Berlin (1997).
 - [16] J. Laherrère and D. Sornette, European Physical Journal B 2, 525-539 (1998).
 - [17] H.Takayasu, M.Takayasu, M.P.Okazaki, K.Marumo and T.Shimizu, Fractal Properties in Economics, in 'Paradigms of Complexity', World Scientific, ed. Miroslav M. Novak, (2000) (ISBN 981-02-4292-1), pp.243-258
 - [18] U. Frisch and D. Sornette, J. Phys. I France 7, 1155 (1997)
 - [19] S. Ghashghaie, W. Breymann, J. Peinke, P. Talkner and Y. Dodge, Nature 381, 767 (1996).
 - [20] A. Arneodo, Muzy, J.F. and Sornette, D., European Physical Journal B 2, 277 (1998)
 - [21] J.-F. Muzy, D. Sornette, J. Delour and A. Arneodo, Multifractal returns and Hierarchical Portfolio Theory, submitted to Quantitative Finance (preprint at <http://arXiv.org/abs/cond-mat/0008069>)
 - [22] R. Friedrich, J. Peinke and C. Renner, Phys. Rev. Lett. 84, 5224 (2000)
 - [23] R. Friedrich and J. Peinke, Phys. Rev. Lett. 78, 863 (1997); A. Naert, R. Friedrich and J. Peinke, Phys. Rev E 56, 6719 (1997).
 - [24] H. Kesten, Acta Math. 131, 207 (1973).

- [25] C. de Calan, Luck, J.-M., Nieuwenhuizen, T.M. & Petritis, D., J. Phys. A 18, 501 (1985).
- [26] D. Sornette, Critical Phenomena in Natural Sciences (Chaos, Fractals, Self-organization and Disorder: Concepts and Tools) (Springer Series in Synergetics, Heidelberg, 2000).
- [27] U.A. Müller, M.M. Dacorogna, R. Dav, R.B. Olsen, O.V. Pictet and J.E. von Weizscker, Volatilities of Different Time Resolutions - Analyzing the Dynamics of Market Components, Journal of Empirical Finance 4, No. 2-3, 213-240 (1997).
- [28] G. Ballocchi, M. M. Dacorogna, R. Gencay, Intraday Statistical Properties of Eurofutures by Barbara Piccinato, Derivatives Quarterly 6 (2), 28-44 (1999).